



HEPATOPROTECTIVE INFLUENCES OF MELATONIN ON THE LEVELS OF ANTIOXIDANTS AND LIPID PEROXIDATION IN HYPERAMMONEMIC WISTAR RATS

Peter Jasmin Lena* and Perumal Subramanian

Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalai Nagar – 608002, Tamil Nadu, India

E-mail: jasminmalligai@gmail.com

*Corresponding author

Published: 20 April 2015

AJBBL 2015, Volume 04: Issue 02 Page 01-08

Received: 17 March 2015

Accepted:

ABSTRACT

The antioxidant potential of melatonin (MLT) on hyperammonemia induced by ammonium acetate treatment was studied in rats. In liver tissue, the levels of thiobarbituric acid reactive substances and lipid profile variables was observed to be increased significantly in ammonium acetate treated rats and decreased significantly in rats treated with melatonin and ammonium acetate. Further, enzymatic, superoxide dismutase, catalase and glutathione peroxidase and non-enzymatic, reduced glutathione antioxidants in liver tissues decreased significantly in ammonium acetate treated rats and increased significantly in rats treated with melatonin and ammonium acetate. These biochemical alterations could be due to the ability of melatonin to (i) scavenge a variety of radicals and reactive oxygen species (ii) induce antioxidative enzymes which reduce steady state levels of reactive oxygen species and (iii) stabilize cell membranes which assist them in reducing oxidative damage and thus could prevent oxidative stress in rats.

INTRODUCTION

Ammonia is a catabolic product of protein and nitrogenous compounds that is formed in mammals and humans. At high levels, ammonia is neurotoxic; it affects the functions of the central nervous system, and leads to coma and death (1). Hyperammonemia, caused by insufficient removal

of ammonia in the liver (2) or portacaval shunting (3), which is responsible for the development of hepatic encephalopathy (4). Ammonia intoxication impairs mitochondrial function (5), which could lead to decreased ATP synthesis and increased formation of free radicals (6). The major toxic



Research Article

effects of ammonia likely involve changes in cellular pH and the depletion of certain citric acid cycle intermediates, in particular α -keto-glutarate. Sustained Hyperammonemia in mice leads to increased lipid peroxidation in liver, reflecting an oxidative stress condition (7, 8). Melatonin (*N*-acetyl-5-methoxy-tryptamine) is the main secretory product of the pineal gland. It is present in virtually all organisms ranging from bacteria (9) to mammals (10). Further, Melatonin is an endogenous free radical scavenger (11) and a broad spectrum antioxidant (12). It detoxifies a variety of free radicals and reactive oxygen intermediates including the hydroxyl radical, peroxy nitrite anion, singlet oxygen and nitric oxide (13). Melatonin, which shows extreme diffusibility through membranes, is important for its scavenging action, since it could enter all cells and every subcellular compartment.

Systematic investigations of the levels of lipid peroxidation products and the levels of enzymatic or non enzymatic antioxidants under the conditions of hyperammonemia are lacking. The present study deals with the levels of thiobarbituric acid reactive substances (TBARS- the products of lipid peroxidation) and the levels of catalase, superoxide dismutase and glutathione peroxidase (enzymatic antioxidants) and reduced glutathione (non-enzymatic antioxidant) in the tissue under the condition levels of hyperammonemia and during melatonin

treatment in rats. Furthermore, the levels of lipid profile variables (free fatty acids, triglycerides, phospholipids and cholesterol) in all the groups were investigated.

MATERIALS AND METHODS

Adult male Wistar rats (180-220g), obtained from National Centre for Laboratory Animal Sciences, Hyderabad, were maintained in polypropylene cages in a controlled environment (22-24°C) under 12:12h light dark cycles. Standard pellet diet purchased from (Kamadhenu Agencies, Bangalore, India) and water were provided *ad libitum*. All studies were conducted in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals (14). Melatonin was purchased from Sisco Research Laboratories Private Limited, Mumbai, India. Ammonium acetate and all other chemicals used in this study were of analytical grade.

The animals were divided into four groups of six rats each and all were fed with the standard pellet diet. Group I animals served as controls. Group II animals were administered with ammonium acetate intraperitoneally (100 mg/kg) every day for 45 days (15). Group III animals were treated with ammonium acetate as group II animals along with melatonin (5mg/kg) intraperitoneally (16). Group IV animals received melatonin (5 mg/kg) intraperitoneally throughout the experiment.



Research Article

The experiment was terminated after 45 days and all animals were killed by cervical decapitation. Blood samples were collected from each group of rats. Rats were sacrificed by cervical dislocation and liver samples were collected and stored at -80°C . Liver samples from each rat were homogenized in freshly prepared phosphate buffer saline. Tissue homogenate was used for the estimation of TBARS (17). Remaining volume of homogenate was centrifuged at 5000g for 15 min at 4°C . The supernatant was collected and used for the estimation of antioxidants such as catalase, superoxide dismutase, glutathione peroxidase and reduced glutathione. The extent of lipid peroxidation (TBARS) in liver was determined by measuring malondialdehyde content based on the reaction with thiobarbituric acid (TBA) (17). Data were expressed nmoles per 100 gm tissue. Catalase activity was determined by measuring the decomposition of hydrogen peroxide at 240nm (18). Data was expressed as units per mg protein. Tissue glutathione content in liver homogenate was measured by biochemical assay using dithionitrobenzoic acid (DTNB) (19). Superoxide Dismutase has been assayed by a spectrophotometric method based on the inhibition of a superoxide – driven NADH oxidation (20). Data was expressed as 50% inhibition of NBT reduction /min/mg protein. Glutathione Peroxidase activity was assayed by following the oxidation of NADH at 340nm in the

presence of glutathione reductase which catalyzed the reduction of GSSG formed by the peroxidase (21). Data was expressed as μg of GPx consumed min/mg protein. Protein concentration was determined by the method of Bradford (22).

Lipids from the tissues were extracted by the method of Folch (23). Total cholesterol was determined by the method of Zlatkis (24). Lipid extract was treated with ferric chloride acetic acid reagent to precipitate the proteins. The protein free supernatant was treated with concentrated sulphuric acid a reddish purple color formed was read in a Spectronic20 colorimeter at 560nm. Values are expressed as mg/100g tissue or mg/dl. Phospholipids were determined by the method of Zilversmit (25). The formation of stable blue color, which was read in a colorimeter at 680nm. The amounts of phospholipids are expressed as mg/100g tissue or mg/dl. Triglycerides were determined by the method of Foster (26). The absorbance of yellow colored compound was read in a Spectronic20 colorimeter at 405nm. The triglyceride content is expressed as mg/100g tissue or mg/dl. Free fatty acids were determined by the method of Falholt (27). Non esterified free fatty acids were estimated by the method of copper soap formation Falholt (27). The amounts of free fatty acids are expressed as mg/100g tissue or mg/dl.



Research Article

The data were analyzed using an analysis of variance (ANOVA) and the group means were compared by Least Significant Difference (LSD) test. The results were considered statistically significant if the p-value was 0.05 or less.

RESULTS AND DISCUSSION

Ammonia is removed either in the form of urea in periportal hepatocyte's and/or as glutamine in perivenous hepatocytes (28). Elevated levels of ammonia, in ammonium acetate treated rats may be

due to the tissue damage caused by ammonia induced free radical generation, leading to oxidative stress and tissue damage. (6,8,30). Melatonin is an effective free radical scavenger (13) which by its antioxidant potential decreases the ammonia levels. Under hyperammonemic conditions, elevated levels of ammonia result in the production of free radicals such as hydroxyl radicals, superoxide radicals, peroxy radicals, alkoxy radicals and reactive nitrogen species.

Table: 1 **Changes in the levels of TBARS and antioxidants in liver**

	TBARS	GSH	SOD	CAT	GPx
Group I	1.77 ± 0.08	23.62 ± 1.97	4.94 ± 0.49	77.13 ± 7.43	12.03 ± 1.24
Group II	3.73 ± 0.32 ^{xxx}	11.30 ± 0.91 ^{xxx}	2.22 ± 0.09 ^{xxx}	34.28 ± 2.35 ^{xxx}	4.71 ± 0.25 ^{xxx}
Group III	2.42 ± 0.16 ^{xxxa}	18.23 ± 0.79 ^{xxx,a}	4.40 ± 0.49 ^{xxx,a}	68.62 ± 3.32 ^{xxxa}	8.78 ± 0.64 ^{xxx,a}
Group IV	1.73 ± 0.10 ^{ns}	23.14 ± 1.61 ^{ns}	5.03 ± 0.51 ^{ns}	74.80 ± 6.06 ^{ns}	12.00 ± 1.16 ^{ns}

Statistical significance was evaluated using ANOVA followed by Least Significant Difference (LSD) test. Group II is compared with Group I (^a p < 0.001), Group III is compared with Group II (p < 0.001), Group IV is compared with Group I; ns not significant. The units for TBARS is n moles/10g tissue, GSH is mg/g tissue, SOD is 50% inhibition of NBT redn /min/mg/protein and CAT is μ moles of H₂O₂ consumed/min/mg/protein. GPx μg of GPx consumed/ min /mg/protein.

Elevated levels of TBARS have been observed in the liver tissue of ammonium acetate treated rats indicating the increased levels of lipid peroxidation (Table 1). It is a well-established fact that ammonia intoxication enhances lipid peroxidation and generates free radicals(6,8,29)

.The levels of TBARS in ammonium acetate and melatonin treated rats were significantly decreased when compared to group 2 rats. This suggests that melatonin could offer protection against lipid peroxidation (30).



Table 2: **Changes of lipid profiles in liver tissue**

	Free fatty acids	Phospholipids	Triglycerides	Cholesterol
Group I	596.76 ± 57.53	950.18 ± 102.82	342.84 ± 29.04	348.62 ± 32.67
Group II	910.75 ± 73.13 ^{xxx}	1618.84 ± 57.37 ^{xxx}	653.06 ± 32.84 ^{xxx}	565.86 ± 55.47 ^{xxx}
Group III	748.06 ± 54.69 ^{xxx,a}	1299.87 ± 98.98 ^{xxx,a}	422.78 ± 30.14 ^{xxx,a}	458.83 ± 33.75 ^{xxx,a}
Group IV	573.30 ± 61.07 ^{ns}	941.01 ± 100.33 ^{ns}	458.83 ± 33.75 ^{xxx,a}	340.62 ± 30.41 ^{ns}

Statistical significance was evaluated using ANOVA followed by Least Significant Difference (LSD) test. Group II is compared with Group I (^a $p < 0.001$). Group III is compared with Group II ($p < 0.001$). Group IV is compared with Group I; ns not significant. Free fatty acids, Phospholipids, Triglycerides, Cholesterol are expressed as mg/100g of tissue.

The non-enzymatic antioxidant glutathione is a scavenger of hydroxyl radicals and singlet oxygen (31). It has been reported that ammonia intoxication induces depletion of glutathione and an increase in lipid peroxidation (5). Reports have also shown that ammonia intoxication leads to the increased formation of nitric oxide which results in the oxidation of glutathione (GSH) to glutathione disulphide (GSSG) and to mixed glutathione disulphides (GSSR) resulting in depletion of GSH and increased free radical formation (32). Group III rats compared to Group II rats showed elevated levels of glutathione. This is because, under hyperammonemic conditions, melatonin increases the levels of glutathione, an important intracellular antioxidant, by stimulating its rate-limiting enzyme, γ -glutamylcysteine synthase (33).

In our study, the decreased activities of antioxidant enzymes (SOD, CAT, GPx) in the ammonium acetate treated group may be due to the inhibition of these enzymes by nitric oxide. It is known that ammonia-induced inhibition of antioxidant enzymes is mediated by the activation of NMDA receptors that leads to increased intracellular calcium levels, which in turn activate neuronal nitric oxide synthase, leading to the formation of nitric oxide which inhibits the activities of antioxidant enzymes (6).

In hyperammonemic conditions, melatonin causes an increase in the gene expression and activities of the antioxidant enzymes such as glutathione peroxidase, glutathione reductase and superoxide dismutase (34,35) which results in the elevated levels of these enzymes in Group II rats. The elevated levels of these enzymes might protect against oxidative damage caused by the free radical formation (36).



Research Article

Melatonin directly scavenges hydrogen peroxide to form N^1 -acetyl- N^2 -forms N^1 -acetyl-5-methoxy Kynuramine (37). These biogenic amines could also scavenge hydroxyl radicals and reduce lipid peroxidation. Ammonium acetate may deplete the levels of α -KG and other Krebs cycle intermediates (38) and thus elevate the levels of acetyl coenzyme A. The elevated levels of acetyl coenzyme A may increase the levels of lipid profile variables (free fatty acids, triglycerides, phospholipids and cholesterol) as observed in our study (Table 2). The decreased α -KG levels in rats treated with ammonium acetate might be

reversed during treatment with melatonin, since melatonin was found to reduce these levels (39).

CONCLUSION

Our results suggest that melatonin could control the oxidative stress caused by hyperammonemia by (i) directly scavenging a variety of radicals and reactive oxygen species (ii) inducing antioxidative enzymes which reduce steady state levels of reactive oxygen species and by (iii) stabilizing cell membranes which assist them in reducing oxidative damage.

REFERENCE

1. PlumF, Hindfelt B, In : Vinken PJ., Bruyn GW., Klawans HI : Eds. Metabolic and deficiency disease of nervous system : the neurological complications of liver disease. New York, Elsevier 1976,349
2. Meijer AJ, Lamers WH, Chamuleau RAFM (1990) Nitrogen metabolism and ornithine cycle function *Physiol Rev* 70:701 – 748.
3. Butterworth RF, Giguere JF, Michaud J, Lavoie J, Pomier-Layrargues G (1987) Ammonia: key factor in the pathogenesis of hepatic encephalopathy. *Neurochem Pathol* 6: 1-12
4. Butterworth RF (1995) Hepatic encephalopathy. *Neurologist* 1: 95-104.
5. Kosenko E., KaminskyY, Kaminsky A.,Valencia M., Lee L. Hermenegildo C., Felipo V Superoxide production and antioxidant enzymes in ammonia intoxication in rats. *Free Radical Res* 1997 27, 637-644.
6. Kosenko E, Kaminsky Y, Stavroskaya I.G and Felipo V.(2000). Alteration of mitochondrial calcium homeostasis by ammonia induced activation of NMDA receptors in rat brain *in vivo*. *Brain Res* 880, 139 – 146.
7. O 'Connor JE, Costell M (1990) New roles of carnitine metabolism in ammonia cytotoxicity In : Grisolia S, Felipo V, Minana MD (eds) : Cirrhosis, hepatic encephalopathy and ammonia toxicity. Plenum press, New York pp 183 – 195



Research Article

8. Dakshayani KB, Velvizhi S, Subramanian P (2002) Effects of ornithine α -ketoglutarate on circulatory antioxidants and lipid peroxidation products in ammonium acetate treated rats. *Ann Nutr Metab* 46: 93-96.
9. Manchester LC, Poeggeler B, Alvares FL, Ogden GB, Reiter RJ (1995) Melatonin immunoreactivity in the photosynthetic prokaryote, *Rhodospirillum rubrum* : implications for an ancient antioxidant system. *Cell Mol Biol Res* 41 :391 – 395.
10. Poeggeler ., B Balzer I., Hardeland R ., Lerchl A., (1991) Pineal hormone melatonin oscillates also in the dinoflagellate *Gonyaulax polyedra*. *Naturwissenschaften* 78, 268 – 269.
11. Tan DX, Chen LD, Poeggeler B, Manchester LC, and Reiter RJ (1993) Melatonin: a potent endogenous hydroxyl radical scavenger *Endocr J.* 1:57-6.
12. Reiter (1993) Interactions of the pineal hormone melatonin as a buffer against macro molecular oxidative damage. *Neurochem Int* 3 : 453 – 460.
13. Reiter RJ, Tan DX, Cabrera J,D' Arpa D, Sainz Rm, Mayo JC, Ramos S (1999) The oxidant/antioxidant network role of melatonin. *Biol Signals Recept* 8:56-63.
14. National Institutes of Health Guide for the Care and Use of Laboratory Animals DHEW publication (NIH), revised (1985) Office of Science and Health Reports, DRR/NIH Bethesda, USA.
15. Hilgier W., Albrecht J., Lisy V., Stastny F (1990). The effect of acute and repeated hyperammonemia on gamma – glutamyl transpeptidase in homogenates and capillaries of various rat brain regions. *Mol Chem Neuropathol* 13: 47 – 56.
16. Liu F, Ng TB (2000) Effect of pineal indoles on the activities of the antioxidant defence enzymes superoxide dismutase, catalase and glutathione reductase and levels of reduced and oxidized glutathione in rat tissues. *Biochem Cell Biol* 78: 447 – 453.
17. Nichans WG, Samuelson B (1968) Formation of malondialdehyde from phospholipid arachidonate during microsomal lipid peroxidation *Eur J Biochem* 6:126 -130.
18. Sinha, KA., Colorimetric assay of catalase. *Anal.Biol.Chem.*, 1972 47, 389 – 394.
19. Ellman GL. (1959) Tissue sulphhydryl groups. *Arch Biochem Biophys* 82: 70 – 79
20. Kakkar P., Das B., Viswanathan DN:(1984) A modified spectrophotometric assay of superoxide dismutase. *IndianJ Biochem. Biophys.*, 21 : 130 – 132.
21. Rotruck JT, Pope AL, Gauthier HE, Swanson AB, Hafeman DG and Hoekstna WG, (1973). Selenium: Biochemical roles as component of glutathione peroxidase. *Science* 178 : 588 – 590.



Research Article

22. Bradford M.M (1976) "Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein – dye binding" Anal Biochem 72 : 248 -254.
23. Folch J, Lees M and Stanley GHS. (1957). A simple method for the isolation and purification of total lipids from animal tissues. J Biol Chem 226: 497- 509.
24. Zlatkis A, Zak B, Boyle AJ (1953) A new method for the direct determination of serum cholesterol. J Lab Clin Med 45:486
25. Zilversmit DB, Mc Chandless EL, Davis AK (1950). Microdetermination of plasma phospholipids by trichloroacetic acid precipitation. J.Lab Clin Med 35:155.
26. Foster CS and Dunn O.(1973). Stable reagents for determination of serum triglycerides by a colorimetric Hantzsch condensation method. Clin Chim Acta 19:338-340.
27. Falholt K, Falholt W, Lund B (1973). An easy colorimetric method for routine determination of free fatty acids in plasma. Clin Chim Acta 19:338.
28. Nelson NL, Cox MM (2000) Lehninghr Principles of Biochemistry Macmillan Worth Publishers, Hampshire,UK
29. Vidya M, Subramanian P (2003) Enhancement of circulatory antioxidants by α - ketoglutarate during sodium valproate treatment in Wistar rats. Pol J Pharmacol 55 : 31 – 36
30. Lastra Ld, Cazeba J, Motiva V, Martin Mj (1997) Melatonin protects against gastric ischemia-reperfusion injury in rats. J Pineal Res 23:47-52.
31. Halliwell B, Gutteridge JMC (1999) Free radicals in biology and medicine 3rd edition. Oxford Science Publications, UK
32. Luperchio S, Tamir S,Tannenbaum SR (1996) NO-induced oxidative stress and glutathione metabolism in rodent and human cells. Free Radical Biol Med 21:513-519
33. Urata Y, Honma S,Goto s, Todoroki S, Ueda T, Cho S, Honma K., Kondo T, (1999) Melatonin induces Y-glutamylcysteine synthase mediated by activator protein-1 in human vascular endothelial cells. Free Radical Biol Med 27: 838-847
34. Barlow-Walden LR,Reiter RJ, Abe M, Pablos M, Pelaez Ma, Chen LD (1995) Melatonin stimulates brain glutathione peroxidase activity. Neurochem Int 26: 497 -502
35. Antolin I, Rodriguez C, Sainz RM (1996) Neurohormone melatonin prevents cell damage effect on gene expression for antioxidant enzymes. FASEB J 10: 882 -890.
36. Reiter RJ and Tan DX (2003). Melatonin: a novel protection agent against oxidative injury of the ischemic reperfused heart. Cardiovasc Res 58:10 – 19.



Research Article

37. Tan DX, Manchester LC, Reiter RJ, Plummer BF, Limson J, Weintraub ST, Qi W (2000) Melatonin directly scavenges H_2O_2 : a potentially new metabolic pathway of melatonin biotransformation. Free Radical Biol Med 29:1177-1185
38. Yamamoto H (1989) Hyperammonemia: increased brain neutral and aromatic amino acid levels and encephalopathy induced by cyanide in mice. J Toxicol Appl Pharmacol 99: 415-420.
39. Baydas G, Yilmaz O, Celik S, Yasar A, Felit Gursu M (2002) Effects of certain micronutrients and melatonin on plasma lipid, lipid peroxidation and homocysteine levels in rats. Arch Med Res 33: 515 – 519.